

Nutritional Deficiency and Its Symptoms:



Nutrient	Incidence of Deficiency	Typical Symptoms and Diseases
Biotin	Uncommon	Dermatitis, eye inflammation, hair loss, loss of muscle control, insomnia, muscle weakness
Calcium	Average diet contains 40 to 50% of RDA*	Brittle nails, cramps, delusions, depression, insomnia, irritability, osteoporosis, palpitations, periodontal disease, rickets, tooth decay
Chromium	90% of diets deficient	Anxiety, fatigue, glucose intolerance, adult-onset diabetes
Copper	75% of diets deficient; average diet contains 50% of RDA*	Anemia, arterial damage, depression, diarrhea, fatigue, fragile bones, hair loss, hyperthyroidism, weakness
Essential fatty acids	Very common	Diarrhea, dry skin and hair, hair loss, immune impairment, infertility, poor wound healing, premenstrual syndrome, acne, eczema, gall stones, liver degeneration
Folic acid	Average diet contains 60% of RDA*; deficient in 100% of elderly & 48% of adolescent girls; requirement doubles in pregnancy	Anemia, apathy, diarrhea, fatigue, headaches, insomnia, loss of appetite, neural tube defects in fetus, paranoia, shortness of breath, weakness
Iodine	Uncommon since introduction of salt with iodine	Cretinism, fatigue, hypothyroidism, weight gain
Iron	Most common mineral deficiency	Anemia, brittle nails, confusion, constipation, depression, dizziness, fatigue, headaches, inflamed tongue, mouth lesions
Magnesium	75 to 85% of diets deficient: average diet contains 50 to 60% of RDA*	Anxiety, confusion, heart attack, hyperactivity, insomnia, nervousness, muscular irritability, restlessness, weakness

Manganese	Unknown, may be common in women	Atherosclerosis, dizziness, elevated cholesterol, glucose intolerance, hearing loss, loss of muscle control, ringing in ears
Niacin	Commonly deficient in elderly	Bad breath, canker sores, confusion, depression, dermatitis, diarrhea, emotional instability, fatigue, irritability, loss of appetite, memory impairment, muscle weakness, nausea, skin eruptions and inflammation
Pantothenic acid (B5)	Average elderly diet contains 60% of RDA*	Abdominal pains, burning feet, depression, eczema, fatigue, hair loss, immune impairment, insomnia, irritability, low blood pressure, muscle spasms, nausea, poor coordination
Potassium	Commonly deficient in elderly	Acne, constipation, depression, edema, excessive water consumption, fatigue, glucose intolerance, high cholesterol levels, insomnia, mental impairment, muscle weakness, nervousness, poor reflexes
Pyridoxine (B6)	71% of male and 90% of female diets deficient	Acne, anemia, arthritis, eye inflammation, depression, dizziness, facial oiliness, fatigue, impaired wound healing, irritability, loss of appetite, loss of hair, mouth lesions, nausea
Riboflavin	Deficient in 30% of elderly Britons	Blurred vision, cataracts, depression, dermatitis, dizziness, hair loss, inflamed eyes, mouth lesions, nervousness, neurological symptoms (numbness, loss of sensation, "electric shock" sensations), seizures. sensitivity to light, sleepiness, weakness
Selenium	Average diet contains 50% of RDA	Growth impairment, high cholesterol levels, increased incidence of cancer, pancreatic insufficiency (inability to secrete adequate amounts of digestive enzymes), immune impairment, liver impairment, male sterility

Thiamin	Commonly deficient in elderly	Confusion, constipation, digestive problems, irritability, loss of appetite, memory loss, nervousness, numbness of hands and feet, pain sensitivity, poor coordination, weakness
Vitamin A	20% of diets deficient	Acne, dry hair, fatigue, growth impairment, insomnia, hyperkeratosis (thickening and roughness of skin), immune impairment, night blindness, weight loss
Vitamin B-12	Serum levels low in 25% of hospital patients	Anemia, constipation, depression, dizziness, fatigue, intestinal disturbances, headaches, irritability, loss of vibration sensation, low stomach acid, mental disturbances, moodiness, mouth lesions, numbness, spinal cord degeneration
Vitamin C	20 to 50% of diets deficient	Bleeding gums, depression, easy bruising, impaired wound healing, irritability, joint pains, loose teeth, malaise, tiredness.
Vitamin D	62% of elderly women's diets deficient	Burning sensation in mouth, diarrhea, insomnia, myopia, nervousness, osteomalacia, osteoporosis, rickets, scalp sweating
Vitamin E	23% of male and 15% of female diets deficient	Gait disturbances, poor reflexes, loss of position sense, loss of vibration sense, shortened red blood cell life
Vitamin K	Deficiency in pregnant women and newborns common	Bleeding disorders

Zinc	68% of diets deficient	Acne, amnesia, apathy, brittle nails, delayed sexual maturity, depression, diarrhea, eczema, fatigue, growth impairment, hair loss, high cholesterol levels, immune impairment, impotence, irritability, lethargy, loss of appetite, loss of sense of taste, low stomach acid, male infertility, memory impairment, night blindness, paranoia, white spots on nails, wound healing impairment
------	------------------------	---

(Source: Total Wellness by Joseph Pizzorno, ND)